

Quantum mutual entropy for two-level ion in a q-analog trap

S. Shelly Sharma

*Depto. de Fisica, Universidade Estadual de Londrina, Londrina, 86040-370,
PR, Brazil*

N. K. Sharma

*Depto. de Matematica, Universidade Estadual de Londrina, Londrina, 86040-370,
PR, Brazil*

Quantum mutual entropy is used as a measure of information content of ionic state due to ion-laser interaction in a q-analog trap. The initial state of the system is a Schrodinger cat state. It is found that the partial mutual entropy is a good measure of the entanglement and purity of the ionic state at $t > 0$.

New insight into quantum mechanical systems has been obtained by applying information theory technics [1–7]. In particular different probability distributions may be compared by using quantum relative entropy [3] and the degree of entanglement measured by quantum mutual entropy [5]. In this article we examine the entropy, quantum relative entropy and quantum mutual entropy for trapped ion- laser system, as a measure of entanglement and quantum information propagation. For an excellent review on quantum relative entropy see Ref. [8].

We consider an ion interacting with a single mode laser in a q-analog harmonic oscillator trap with the system Hamiltonian given by

$$H_q = \frac{1}{2}\hbar\omega(AA^\dagger + A^\dagger A) + \frac{1}{2}\hbar\Delta\sigma_z + \frac{1}{2}\hbar\Omega(F_q\sigma^+ + F_q^\dagger\sigma^-), \quad (1)$$

where $A^\dagger$ and A are the creation and annihilation operators for the trap quanta satisfying the commutation relations,

$$AA^\dagger - qA^\dagger A = q^{-N} \quad ; \quad NA^\dagger - A^\dagger N = A^\dagger \quad ; \quad NA - AN = -A, \quad (2)$$

q the trap deformation parameter ($q = \exp(\tau)$ with τ real), $\Delta = \omega_a - \omega_l$ the detuning parameter, and Ω the Rabi frequency. The pseudo-spin operators of the two-level ion, $\sigma^\pm$ and σ_z operate on state vectors $|g\rangle$ and $|e\rangle$ where indices g and e stand for the ground and excited states of the two level ion. The operator $F_q = e^{\left(\frac{-|e|^2}{2}\right)} e^{i\epsilon A^\dagger} e^{i\epsilon A}$, and N is the number operator for trap quanta, .

Treating internal and motional degrees of freedom as separate systems, the entropies are defined through the corresponding reduced density operators by

$$S_{ion(c.m.)} = -tr_{ion(c.m.)} \{ \rho_{ion(c.m.)} \ln(\rho_{ion(c.m.)}) \} \quad (3)$$

where the reduced density operators are

$$\rho_{ion(c.m.)} = tr_{c.m.(ion)} \{ \rho \}. \quad (4)$$

We consider the initial state of the trapped ion to be a pure state with

$$\rho_0^{(1)} = |g\rangle \langle g| \otimes |\beta\rangle_q \langle \beta|, \text{ or } \rho_0^{(2)} = |e\rangle \langle e| \otimes |-\beta\rangle_q \langle -\beta|, \quad (5)$$

where the motional state of the ionic center of mass is a q - analog coherent state

$$|\beta\rangle_q = \frac{1}{\sqrt{\exp_q(|\beta|^2)}} \sum_{n=0}^{\infty} \frac{\beta^n}{\sqrt{[n]_q!}} |n\rangle_q. \quad (6)$$

A pure state at $t = 0$ evolves into an entangled state for $t > 0$ with the Hamiltonian H_q governing the dynamics of the system through

$$H_q \Psi^{(i)}(t) = i\hbar \frac{d}{dt} \Psi^{(i)}(t) \quad (7)$$

($i = 1$ for ionic state $|g\rangle$ and for ionic state $|e\rangle$).

Working in the space spanned by basis vectors $\{|u_n\rangle\} \equiv |g, m\rangle, |e, m\rangle, m = 0, 1, \dots, \infty$, the state of the system may be written as

$$\Psi^{(i)}(t) = \sum_m g_m^{(i)}(t) |g, m\rangle + \sum_m e_m^{(i)}(t) |e, m\rangle \quad . \quad (8)$$

We next define the populations and coherences for the system prepared in initial state $|i, \beta\rangle_q$, with the populations defined as

$$P_g^{(i)}(t) = \langle g | \text{tr}_{ion} \rho^{(i)}(t) | g \rangle = \sum_n |g_n^{(i)}(t)|^2 \quad (9)$$

and

$$P_e^{(i)}(t) = \langle e | \text{tr}_{ion} \rho^{(i)}(t) | e \rangle = \sum_n |e_n^{(i)}(t)|^2 \quad (10)$$

The coherences for this initial states are given by

$$C_{ge}^{(i)}(t) = \langle g | \text{tr}_{ion} \rho^{(i)}(t) | e \rangle = \sum_n \left(g_n^{(i)}(t) \right)^* e_n^{(i)}(t) \quad . \quad (11)$$

and

$$C_{eg}^{(i)}(t) = \langle e | \text{tr}_{ion} \rho^{(i)}(t) | g \rangle = \left(C_{ge}^{(i)}(t) \right)^* \quad (12)$$

A straightforward calculation shows that

$$\begin{aligned} S_{ion}^{(i)} = & - \left[P_g^{(i)}(t) \log P_g^{(i)}(t) + P_e^{(i)} \log P_e^{(i)}(t) \right. \\ & \left. + 2 * \text{Re } C_{ge}^{(i)}(t) \log \left(C_{ge}^{(i)}(t) \right)^* \right] \quad . \end{aligned} \quad (13)$$

As at $t = 0$, ionic state and center of mass motional state are pure states, the total entropy of the system is zero. At $t > 0$ the entropies of the center of mass motion and the ionic subsystems are precisely equal. The entropy S_{ion} is a good a measure of entanglement of the system and has more information content than the population inversion which is an experimentally measurable property of the system.

Next we consider trapped ion prepared in the initial state $\Psi(t = 0) = \frac{|g, \beta\rangle_q + |e, -\beta\rangle_q}{\sqrt{2}}$, which is a Macroscopic superposition of states. This is not a pure state but in the phase space the components $|g, \beta\rangle_q$ and $|e, -\beta\rangle_q$ are well separated and distinguishable. The density operator for the internal states of the ion at $t = 0$ is

$$\rho_{ion}(t = 0) = \frac{1}{2} (|g\rangle \langle g| + |e\rangle \langle e|) \quad (14)$$

The state of the system at a time t given

$$\Psi(t) = \sum_m g_m(t) |g, m\rangle + \sum_m e_m(t) |e, m\rangle \quad (15)$$

is a solution of the time dependent Schrodinger equation

$$H_q \Psi(t) = i\hbar \frac{d}{dt} \Psi(t) \quad (16)$$

Defining the probabilities of finding the ion in the ground state and the excited state (the populations) to be respectively

$$P_g(t) = \langle g | \text{tr}_{ion} \rho(t) | g \rangle = \sum_n |g_n(t)|^2 \quad (17)$$

and

$$P_e(t) = \langle e | tr_{ion} \rho(t) | e \rangle = \sum_n |e_n(t)|^2. \quad (18)$$

and the non-diagonal matrix element of the operator $tr_{ion} \rho(t)$ (coherence),

$$C_{ge} = \langle g | tr_{ion} \rho(t) | e \rangle = \sum_n g_n^*(t) e_n(t). \quad (19)$$

We find that the entropy

$$S_{ion} = - [P_g(t) \log P_g(t) + P_e \log P_e(t) + 2 * \text{Re } C_{ge}(t) \log (C_{ge}(t))^*] . \quad (20)$$

We plot in figure I(a) $Sg = S_{ion}^{(1)}$ for $\tau = 0.0$ (Harmonic oscillator trap) , figure I(b) $Scat = S_{ion}$ for $\tau = 0.0$, and figure I(c) $Scat = S_{ion}$ for $\tau = 0.004$ (q -deformed harmonic oscillator trap) as a function of rescaled time parameter $t(\frac{\Omega t}{2\pi})$ for $\beta = 4$. The parameter values used in the numerical calculation are $\overline{\omega} = \frac{\omega}{\Omega} = 50$, $\overline{\epsilon} = \frac{\epsilon}{\Omega} = 0.05$, and $\overline{\Delta} = \frac{\Delta}{\Omega} = -50$. The maximum value of n in Eq. (6) is restricted to $n = 32$. For the case (a) where initial state of the ion is a pure state with $Sg = 0$, the value of Sg increases with t indicating entanglement. After an interval of time characteristic of the system Sg is found to go to zero again indicating a decoupling of the internal and motional degrees of freedom of the ion. In case (b) $Scat$ follows more or less the population inversion. For the case (c) no conclusive information can be obtained about the state of the system at hand.

In order to measure the quantum interference effects due to the initial state superposition of component states $|g, \beta\rangle_q$ and $|e, -\beta\rangle_q$ we use the notion of quantum mutual entropy. Furuichi and co-workers [9] have applied the concept of quantum mutual entropy to dynamical change of state of the atom in Jaynes-Cummings model(JCM) [10]. Quantum relative entropy [7,1] between the two states σ and ρ defined as

$$S(\sigma, \rho) = Tr \sigma (\ln \sigma - \ln \rho) , \quad (21)$$

is a measure of difficulty of distinguishing between the states σ and ρ . The correlation between the initial state $\rho_{ion}(t=0)$ having component states $|i\rangle \langle i|$ ($i = 1, 2$) and the state $\rho_{ion}(t)$ at $t > 0$ can be measured by quantum relative entropies

$$S(\sigma_1^{(i)}, \sigma_2) = tr \sigma_1^{(i)} (\ln \sigma_1^{(i)} - \ln \sigma_2) ; (i = 1, 2) , \text{ where the states} \quad (22)$$

$$\sigma_1^{(i)} = \frac{1}{2} |i\rangle \langle i| \otimes \rho_{ion}^{(i)}(t) \text{ and } \sigma_2 = \rho_{ion}(t=0) \otimes \rho_{ion}(t). \quad (23)$$

The quantum mutual entropy

$$\begin{aligned} I(\rho_{ion}(t=0), \rho_{ion}(t)) &= \sum_{i=1,2} \frac{1}{2} S(\sigma_1^{(i)}, \sigma_2) \\ &= \sum_{i=1,2} \frac{1}{2} \sum_{k=1,2} \left[\langle g | \sigma_1^{(i)} | k \rangle \left(\ln \frac{\langle k | \sigma_1^{(i)} | g \rangle}{\langle k | \sigma_2 | g \rangle} \right) \right. \\ &\quad \left. + \langle e | \sigma_1^{(i)} | k \rangle \left(\ln \frac{\langle k | \sigma_1^{(i)} | e \rangle}{\langle k | \sigma_2 | e \rangle} \right) \right] . \end{aligned} \quad (24)$$

measures the degree of entanglement of the state $\rho_{ion}(t)$. We can verify that in terms of the populations and the coherences defined above the relative quantum entropies for the system starting in initial state given by Eq. (14) are

$$\begin{aligned} S(\sigma_1^{(i)}, \sigma_2) &= \left[P_g^{(i)}(t) \log \left(\frac{P_g^{(i)}(t)}{P_g(t)} \right) + P_e^{(i)} \log \left(\frac{P_e^{(i)}(t)}{P_e(t)} \right) \right. \\ &\quad \left. + 2 * \text{Re} \left[C_{ge}^{(i)}(t) \log \left(\frac{C_{ge}^{(i)}(t)}{C_{ge}(t)} \right)^* \right] \right] . \end{aligned} \quad (25)$$

Substituting Eq. (25) in the definition of quantum mutual entropy for the system, we have

$$I(\rho_{ion}(t=0), \rho_{ion}(t)) = \sum_{i=1,2} \frac{1}{2} \left[P_g^{(i)}(t) \log \left(\frac{P_g^{(i)}(t)}{P_g(t)} \right) + P_e^{(i)} \log \left(\frac{P_e^{(i)}(t)}{P_e(t)} \right) \right. \\ \left. + 2 * \text{Re} \left[C_{ge}^{(i)}(t) \log \left(\frac{C_{ge}^{(i)}(t)}{C_{ge}(t)} \right)^* \right] \right] . \quad (26)$$

We can split $I(\rho_{ion}(t=0), \rho_{ion}(t))$ in to two distinct parts, one depending on populations and the other on off diagonal coherences that is

$$I(\rho_{ion}(t=0), \rho_{ion}(t)) = S(P) + S(C) \quad (27)$$

where

$$S(P) = \sum_{i=1,2} \lambda(i) \left[P_g^{(i)}(t) \log \left(\frac{P_g^{(i)}(t)}{P_g(t)} \right) + P_e^{(i)}(t) \log \left(\frac{P_e^{(i)}(t)}{P_e(t)} \right) \right] . \quad (28)$$

We define population inversion, the difference between the probabilities of finding the system in the ground state and the excited state, as

$$Inv = P_g(t) - P_e(t),$$

which is the most commonly used measure of entanglement for ion in trap system.

Figure II is a plot of quantum mutual entropy I , partial quantum mutual entropy $S(P)$ and population inversion Inv as a function of rescaled time parameter $t(\frac{\Omega t}{2\pi})$ for $\beta = 4$. The q -deformed oscillator trap is characterized by deformation parameter $\tau = 0.004$ ($q = \exp(\tau)$ with τ real). This particular system, studied in refs. [11,12] develops extremely well defined collapses and revivals [13,14] of population inversion as is evident from figure II(a). The quantum mutual entropy I measures the system correlations at collapse, revival peaks and during the transition period when the revival is building up or decaying. As expected the transition periods correspond to highly correlated system. The partial quantum mutual entropy $S(P)$ containing no contributions from coherences shows a very regular pattern with peaks characterizing each collapse and revival. The onset of each collapse and revival is characterized by $S(P) = 0$ and fluctuating I value. The $S(p)$ peaks represent the t values for which the information content of the ionic system is the maximum. With time quantum mutual entropy I as well as $S(P)$ are seen to decrease indicating a dissipative irreversible change in the ionic state. In ref. [9], where quantum mutual entropy dynamics for Jaynes-Cummings model has been reported the coherences turn out to be zero ($S(P) = I$). As such our results for partial mutual quantum entropy $S(P)$ are similar to the results for total quantum mutual entropy I of ref. [9]. From the location of $S(P)$ peaks the successive Inv collapses occur at $t = 67.8, 201.2$, and 336.8 while the revival peaks are seen at $t = 133.2$ and 266.6 .

Acknowledgments

S. S. S. and N. K. S. would like to acknowledge financial support from Universidade Estadual de Londrina, Brazil.

-
- [1] M. Ohya and P. Petz, "Quantum Entropy and Its Use", Texts and Monographs in Physics, (Berlin: Springer-Verlag) (1993).
 - [2] Bennett, D. P. DiVincenzo, Smolin, J. A. and Wootters, W. K. , Phys. Rev. A, **54**, 3824 (1996).
 - [3] V. M. Vedral, B. Plenio, M. A. Rippin and P. L. Knight, Phys. Rev. Lett., **78**, 2275 (1997).
 - [4] C. P. Panos and S. E. Massen, Int. J. of Mod. Phys. E **6** , 497 (1997)
 - [5] V. P. Belavkin and M. Ohya, quant-ph/9812082
 - [6] M. Ohya, Rep. Math. Phys. **27**, **19** (1989)
 - [7] H. Umegaki, Kodai Math. Sem. Rep. **14**, 59(1962).
 - [8] V. Vedral, Preprint quant-ph/0102094 .
 - [9] S. Furuichi, M. Ohya, H. Suyari, Rept. Math. Phys. **44**, 81-86(1999).
 - [10] Jaynes E. T. and Cummings F. W., Proc. Inst. Electr. Eng., **51**, 89 (1963).

- [11] S. Shelly Sharma, N. K. Sharma and Larry Zamick, Phys. Rev. A **56**, 694 (1997) and preprint quant-ph/9506036.
- [12] S. Shelly Sharma and N. K. Sharma, Preprint quant-ph/0011090 .
- [13] F. W. Cummings, Phys. Rev., **140**, A1051 (1965)
- [14] B. W. Shore and P. L. Knight, J. Mod. Optics, **40**, 1195 (1993) .




FIG. 1. (a) $Sg = S_{ion}^{(1)}$ for $\tau = 0.0$ (Harmonic oscillator trap), (b) $Scat = S_{ion}$ for $\tau = 0.0$, and (c) $Scat = S_{ion}$ for $\tau = 0.004$ (q -deformed harmonic oscillator trap) as a function of rescaled time parameter $t(\frac{\Omega t}{2\pi})$ for $\beta = 4$.




FIG. 2. Quantum mutual entropy I , partial quantum mutual entropy $S(P)$ and population inversion Inv as a function of rescaled time parameter $t(\frac{\Omega t}{2\pi})$ for $\beta = 4$ and $\tau = 0.004$.